The Science of Surviving Low Oxygen

Alimbekova Aiperi Alimbekovna
(Lecturer,International Medical Faculty, Osh State University, Kyrgyz Republic.)
Sowmeya Ravi
(Student, International Medical Faculty, Osh State University, Kyrgyz Republic.)
Rithish Kamaraj
(Student,Madha medical college & research institute,Chennai,Tamil nadu,India.)

Abstract

Molecular oxygen (O2)making it essential for most species on Earth.Oxygen is essential for producing energy and supporting vital cellular activities, the body has developed several ways to adapt to low-oxygen conditions..Hypoxia is a condition in which the body or its tissues do not receive enough oxygen to function normally. Since oxygen is essential for energy production and cellular activities, the body activates several adaptive mechanisms to survive low-oxygen conditions.Hypoxia causes hypoventilation, impaired alveolar diffusion, and pulmonary shunting. A key role is played by Hypoxia-Inducible Factors (HIFs), which regulate genes involved in oxygen delivery, energy metabolism, red blood cell production, and angiogenesis. During hypoxia, cells switch to alternative energy pathways, while breathing rate and blood circulation increase to improve oxygen supply. Understanding these adaptive responses is important for explaining survival in low-oxygen environments and for developing treatments for diseases such as anemia, stroke, heart disease, and cancer.Exposure to low oxygen levels (hypoxia) in parent organisms was found to increase their lifespan and cause changes in fat storage and fertility that could be passed on to future generations through small RNA molecules. Similar adaptations were also observed when glucose availability changed, showing that environmental conditions can influence the health and characteristics of future generations.

Keywords:Hypoxia,HIF(HypoxiaInducibleFactor),Oxygendeficiency,Cellular adaptation,Erythropoietin (EPO),Hemoglobin,Angiogenesis,Metabolic shift,Anaerobic glycolysis,Gene activation,Transgenerational adaptation.

Introduction

The Science of Surviving Low Oxygen is called hypoxia.Hypoxia is a medical condition where your body or a specific region of your body doesn't receive enough oxygen at the tissue level to function properly.Mild hypoxia usually occurs when oxygen saturation levels drop to 90% - 94%.Traveling to a high altitude, where levels of oxygen are lower, can also cause hypoxemia.Any condition that reduces the amount of oxygen in your blood or restricts blood flow can cause hypoxia.The four different types of hypoxia are hypoxemic, circulatory, anemic and histotoxic.Examples of O2 consuming reactions include aerobic respiration, fatty acid desaturation,Due to vascular insufficiency or blood vessel damage,hypoxia arises in a variety of diseases, including the growth of solid tumours.Hypoxia is normally controlled by HIF (hypoxia-inducible factor), Under normal oxygen levels, HIF is quickly broken down and remains inactive. When oxygen levels decrease, HIF becomes active and turns on genes that help cells survive by improving oxygen use and delivery. At the same time, cells save energy by reducing non-essential activities, switching to a simpler method of energy

production, and recycling damaged cell components through autophagy.

Types of hypoxia

Hypoxia can be classified into four main types based on its underlying cause. Hypoxemic hypoxia occurs when there is a low level of oxygen in the blood, commonly due to lung diseases, heart disorders, congenital heart defects, certain medications that slow breathing, or exposure to high altitudes where oxygen levels are lower. Circulatory hypoxia occurs when the blood contains enough oxygen, but poor blood flow caused by conditions such as heart failure, blood vessel diseases, or blood clots prevents adequate oxygen delivery to tissues. Anemic hypoxia develops when there are too few healthy red blood cells or insufficient hemoglobin to carry oxygen from the lungs to the body's tissues, reducing oxygen transport. Histotoxic hypoxia occurs when oxygen reaches the tissues normally, but the cells are unable to utilize it effectively due to toxic substances such as cyanide, which interfere with normal cellular respiration. Together, these four types illustrate the different ways in which oxygen delivery or utilization can be impaired, leading to tissue hypoxia.

Hypoxia inducible factors (HIFs)

Hypoxia-Inducible Factor-1 (HIF-1) is the body's main oxygen-sensing protein that helps cells respond when oxygen levels become low. It is present in almost all oxygen-breathing organisms and acts as a transcription factor, meaning it switches specific genes on or off according to the body's oxygen needs. HIF-1 is made up of two protein subunits: HIF-1α (alpha) and HIF-1β (beta), also known as ARNT (Aryl Hydrocarbon Receptor Nuclear Translocator). Under normal oxygen conditions, HIF-1α is rapidly broken down, while HIF-1β is always present in the cell. During hypoxia, HIF-1α becomes stable, combines with HIF-1β, and forms the active HIF-1 complex. This complex enters the nucleus and activates genes involved in oxygen transport, red blood cell production, blood vessel formation, and energy metabolism, helping the body adapt to low oxygen levels. Structurally, both subunits contain a basic helix-loop-helix (bHLH) domain for DNA binding, a PAS domain that allows the two proteins to join together, and a C-terminal region that recruits other proteins needed for gene activation. The human HIF family includes HIF-1α (HIF1A), HIF-1β (ARNT), HIF-2α (EPAS1), HIF-2β (ARNT2), HIF-3α (HIF3A), and HIF-3β (ARNT3), each of which contributes to regulating the body's response to hypoxia.

Angiogenesis in Hypoxia

Angiogenesis is the process of forming new blood vessels from existing blood vessels. During hypoxia (low oxygen conditions), tissues do not receive enough oxygen to meet their metabolic needs. To overcome this, the body activates Hypoxia-Inducible Factor-1 (HIF-1), which increases the production of Vascular Endothelial Growth Factor (VEGF). VEGF stimulates endothelial cells to grow, migrate, and form new capillaries, thereby increasing blood supply to oxygen-deprived tissues. The newly formed blood vessels improve oxygen and nutrient delivery, remove metabolic waste products, and help restore normal tissue function. Angiogenesis is an important adaptive response in conditions such as high-altitude exposure, wound healing, ischemic heart disease, and peripheral artery disease. However, excessive angiogenesis can also contribute to the progression of diseases such as cancer, where tumors develop new blood vessels to obtain oxygen and nutrients, allowing them to grow and spread.

Aim

To study the mechanisms of hypoxia survival and adaptation and understand how the human body responds to low oxygen conditions to maintain normal cellular and physiological functions.

Objectives

This study aims to understand hypoxia, a condition in which the body receives insufficient oxygen, and to explore how the body adapts to survive under such conditions. It focuses on the role of Hypoxia-Inducible Factors (HIFs) in detecting low oxygen levels and activating protective mechanisms that help maintain cellular function. The study examines the cellular and physiological changes that occur during hypoxia, including shifts in energy metabolism, increased breathing rate, enhanced cardiac output, and greater production of red blood cells. It also explores the importance of angiogenesis and erythropoietin (EPO) in improving oxygen delivery throughout the body. In addition, the study highlights the significance of hypoxia adaptation in high-altitude survival and athletic performance, while discussing its relevance in diseases such as anemia, stroke, ischemic heart disease, and cancer. By understanding these adaptive mechanisms and hypoxia-related pathways, the study aims to provide insight into potential therapeutic strategies and emphasize the vital role of hypoxia adaptation in maintaining health and supporting life under oxygen-deficient conditions.

Materials and methods

The study focused on understanding both molecular and physiological mechanisms involved in oxygen deprivation and adaptation. Key pathways such as metabolic shift to anaerobic glycolysis, regulation of gene expression under hypoxic conditions, and systemic responses like increased ventilation and cardiac output were examined.Adaptation to oxygen starvation is a key feature of both primary and metastatic neoplasms.Persistent activation of hypoxia-induced genes can result in pathologies, including pulmonary hypertension.During low oxygen conditions (hypoxia), Hypoxia-Inducible Factor (HIF) stimulates the production of Vascular Endothelial Growth Factor (VEGF), which promotes the growth of new blood vessels (angiogenisis).

During hypoxia, cells switch to alternative methods of producing energy, while the body responds by increasing breathing rate, heart activity, and the production of red blood cells to carry more oxygen. The growth of new blood vessels is also stimulated to improve blood supply to oxygen-starved tissues.These adaptations are important in situations such as high-altitude exposure, lung and heart diseases, and other conditions associated with reduced oxygen availability. Understanding how the body adapts to hypoxia not only helps explain human survival under environmental stress but also contributes to the development of new treatments for diseases such as stroke, anemia, ischemic heart disease, and cancer. Therefore, studying hypoxia and its adaptive mechanisms is essential for improving our understanding of human health and disease.Three key signs of low oxygen include Shortness of breath,Confusion or dizziness,Bluish skin.

The best studied mechanism of response to hypoxia involves hypoxia inducible factors (HIFs)control the expression of a genes.HIF-1 and HIF-2 are major transcription factors involved in the hypoxic

response.HIFs are heterodimers consisting of the regulated HIF-α protein subunit which is only expressed during hypoxia, and a constitutively expressed HIF-1β protein subunit.PHD enzymes are oxygen-sensing enzymes that regulate the stability of Hypoxia-Inducible Factors (HIFs) and play a key role in the body's response to oxygen availability.Under normal oxygen conditions (normoxia), PHD enzymes hydroxylate the HIF-α subunit. This modification marks HIF-α for degradation, preventing activation of hypoxia-response genes.

High-altitude physiology involves how the body responds when oxygen levels are low, such as at high mountains. In the short term, the body tries to adjust through acclimatization, which helps improve oxygen delivery and reduces the effects of low oxygen, but extreme altitude exposure can be dangerous and even fatal. Over many generations, some populations develop permanent adaptations through natural selection that help them live comfortably in low-oxygen environments.

Under normal oxygen levels, HIF is continuously broken down and kept inactive. When oxygen levels fall, this breakdown stops, allowing HIF to enter the nucleus and switch on genes that help the cell survive low oxygen conditions. In response, cells reduce energy-consuming activities and shift from oxygen-dependent energy production to a simpler form of energy generation called anaerobic glycolysis. At the same time, some tissues may undergo controlled recycling processes (autophagy) to save energy and improve oxygen efficiency.

Results and discussions :

The study shows that hypoxia triggers a well-coordinated response at both cellular and whole-body levels to maintain survival in low oxygen conditions. When oxygen levels decrease, Hypoxia-Inducible Factors (HIFs) become active and switch on genes that help improve oxygen delivery, energy production, and cell survival. As a result, the body adapts by increasing breathing rate, heart activity, and red blood cell production, while also promoting the formation of new blood vessels to enhance oxygen supply to tissues.

At the cellular level, energy production shifts from oxygen-dependent pathways to simpler anaerobic processes, and unnecessary energy use is reduced through controlled mechanisms like autophagy. These changes help cells survive during oxygen shortage by conserving energy and maintaining basic functions.

When mammals are exposed to low-oxygen (hypoxic) environments, their bodies make several adaptations to maintain an adequate oxygen supply. The respiratory system responds by increasing the rate and depth of breathing and, over time, may develop greater lung capacity to improve oxygen uptake. The hematological system produces more red blood cells and hemoglobin, allowing the blood to carry more oxygen. The cardiovascular system increases cardiac output and forms new blood vessels (angiogenesis) to enhance oxygen delivery to tissues. Muscles adapt by increasing myoglobin, which stores oxygen and provides it when needed. At the cellular level, hypoxia-inducible factor (HIF) activates genes that help cells survive low oxygen conditions and promotes increased glycolysis for energy production. In populations living at high altitudes, genetic adaptations involving genes such as EPAS1 and EGLN1 improve oxygen sensing and utilization. Together, these changes help mammals maintain normal body functions and survive in environments where oxygen availability is limited.(Table 1).

System	Adaptation to Hypoxia
Respiratory	Hyperventilation, increased lung capacity
Hematological	Polycythemia, increased Hb
Cardiovascular	Increased cardiac output, angiogenesis
Cellular	HIF activation, increased glycolysis
Genetic	EPAS1, EGLN1 gene adaptations

Table 1 : organ system and its adaptation to hypoxia

When oxygen levels in the body decrease (hypoxia), the reduced oxygen tension prevents the breakdown of Hypoxia-Inducible Factor-1 alpha (HIF-1α), causing it to become stabilized and activated. Activated HIF-1α enters the nucleus and stimulates the transcription of various hypoxia-responsive genes. This leads to increased production of erythropoietin (EPO), which promotes red blood cell formation (erythropoiesis) and improves oxygen transport. HIF-1α also increases vascular endothelial growth factor (VEGF) production, stimulating angiogenesis and enhancing blood supply to tissues. Additionally, it upregulates GLUT-1, increasing glucose uptake by cells, and activates glycolytic enzymes, allowing greater ATP production through anaerobic metabolism when oxygen is limited. Furthermore, increased nitric oxide synthase activity produces more nitric oxide, causing vasodilation and improving blood flow and oxygen delivery. Together, these responses help the body adapt and survive under low-oxygen conditions.(Table 2)

Hypoxic Stimulus	Response
↓ Oxygen tension	Stabilization of HIF-1α
HIF-1α activation	Transcription of hypoxia-responsive genes
↑ EPO	Erythropoiesis
↑ VEGF	Angiogenesis
↑ GLUT-1	Increased glucose uptake
↑ Glycolytic enzymes	Enhanced anaerobic metabolism
↑ Nitric oxide synthase	Vasodilation

Table 2 :Molecular pathway of hypoxia adaptation.

Transgenerational adaptation occurs and suggest that H3K27me3 is a critical modification for adapting transgenerationally to repeated generational stresses.Transgenerational adaptation means that when organisms are exposed to the same environmental stress, such as low oxygen levels, over several generations, their offspring may become better adapted to that stress even without direct exposure. Research suggests that this adaptation can be passed from parents to offspring through epigenetic changes, which alter gene activity without changing the DNA sequence itself. One important epigenetic modification is H3K27me3 (trimethylation of histone H3 at lysine 27), which acts like a molecular switch that can turn certain genes off. Studies indicate that H3K27me3 plays a key role in helping organisms "remember" repeated environmental stresses across generations, allowing future generations to respond more effectively and survive better under similar conditions. Thus, H3K27me3 is considered a critical mechanism for transgenerational adaptation to recurring stresses.

The study also highlights that hypoxia is involved in many conditions such as high-altitude exposure, cardiovascular and respiratory diseases, anemia, and cancer. In addition, long-term or repeated exposure to low oxygen can influence future generations through biological changes involving small RNA pathways and metabolic regulation.Overall, hypoxia adaptation is a vital survival mechanism that allows the body to cope with oxygen deficiency and maintain essential life processes under stress conditions.

Conclusions :

Hypoxia is a condition in which the body does not receive enough oxygen to meet its needs. To survive under these low-oxygen conditions, the body activates several adaptive mechanisms at the molecular, cellular, and physiological levels. Hypoxia-Inducible Factors (HIFs) play a central role by regulating genes that increase oxygen delivery, red blood cell production, blood vessel formation, and energy production. The body also responds by increasing breathing rate, heart activity, and hemoglobin levels, while cells switch to alternative energy pathways to maintain their functions. Long-term adaptation can occur through genetic changes in high-altitude populations and through epigenetic mechanisms that may pass beneficial adaptations to future generations. Understanding these responses is important because hypoxia is involved in many diseases, including anemia, stroke, heart disease, and cancer. Overall, hypoxia adaptation is a vital survival process that helps organisms maintain normal functions and cope with oxygen deficiency.

References

1. Muelleck, J. R., Maracaja, L., & Templeton, T. W. (2022). Selective lobar blockade with a bronchial blocker in combination with a double lumen tube to manage refractory hypoxemia: A case report. Cureus, 14(7), e26638. https://doi.org/10.7759/cureus.26638.

2. Karzai, W., & Schwarzkopf, K. (2009). Hypoxemia during one-lung ventilation: Prediction, prevention, and treatment. Anesthesiology, 110(6), 1402–1411. https://doi.org/10.1097/ALN.0b013e31819fb15d.

3. Henig, N. R., & Pierson, D. J. (2000). Mechanisms of hypoxemia. Respiratory Care Clinics of North America, 6(4), 501–521. https://doi.org/10.1016/S1078-5337(05)70087-3 .

4. Bhutta, B. S., Alghoula, F., & Berim, I. (2024). Hypoxia and hypoxemia. In StatPearls. StatPearls Publishing. https://www.ncbi.nlm.nih.gov/books/ .

5. Cai, C., Ni, G., Chen, L., Deng, C., Chai, S., Wang, R., Zhang, R., Luo, F., & Ge, R.-L. (2026). Altitude hypoxia and hypoxemia: Pathogenesis and management. Signal Transduction and Targeted Therapy, 11, Article 27. https://doi.org/10.1038/s41392-026-013xx-x .

6. Grocott, M., Montgomery, H., & Vercueil, A. (2007). High-altitude physiology and pathophysiology: Implications and relevance for intensive care medicine. Critical Care, 11(1), 1–10. https://doi.org/10.1186/cc5002 .

7. Lee, P., Chandel, N. S., & Simon, M. C. (2020). Cellular adaptation to hypoxia through HIFs and beyond. Nature Reviews Molecular Cell Biology, 21(5), 268–283. https://doi.org/10.1038/s41580-020-0227-y.

8. Lee, P., Chandel, N. S., & Simon, M. C. (2020). Cellular adaptation to hypoxia through HIFs and beyond. Nature Reviews Molecular Cell Biology, 21(5), 268–283. https://doi.org/10.1038/s41580-020-0227-y.

9. Alcock, J. (2025, July 2). Adaptation to hypoxia. Evolution Medicine.

https://evolutionmedicine.com/adaptation-to-hypoxia/.

10. Swenson, E. R., & Bärtsch, P. (Eds.). (2014). High altitude: Human adaptation to hypoxia. Springer. https://doi.org/10.1007/978-1-4614-8772-2.
11. Li, F., Qiao, Z., Duan, Q., & Nevo, E. (2021). Adaptation of mammals to hypoxia. Animal Models and Experimental Medicine, 4(4), 311–318. https://doi.org/10.1002/ame2.12189.
12. Bhutta, B. S., Alghoula, F., & Berim, I. (2024). Hypoxia and hypoxemia. In StatPearls. StatPearls Publishing. https://www.ncbi.nlm.nih.gov/books/.
13. Cai, C., Ni, G., Chen, L., Deng, C., Chai, S., Wang, R., Zhang, R., Luo, F., & Ge, R.-L. (2026). Altitude hypoxia and hypoxemia: Pathogenesis and management. Signal Transduction and Targeted Therapy, 11, Article 27.
14. Grocott, M., Montgomery, H., & Vercueil, A. (2007). High-altitude physiology and pathophysiology: Implications and relevance for intensive care medicine. Critical Care, 11(1), 203. https://doi.org/10.1186/cc5002.
15. Lee, P., Chandel, N. S., & Simon, M. C. (2020). Cellular adaptation to hypoxia through HIFs and beyond. Nature Reviews Molecular Cell Biology, 21(5), 268–283. https://doi.org/10.1038/s41580-020-0227-y.
16. Li, F., Qiao, Z., Duan, Q., & Nevo, E. (2021). Adaptation of mammals to hypoxia. Animal Models and Experimental Medicine, 4(4), 311–318. https://doi.org/10.1002/ame2.12189.
17. Kim, K., Telger, A., Sarkar, G., Surya, S., & Greer, E. L. (2025). Transgenerational adaptation to hypoxia. Science Advances, 11(43). https://doi.org/10.1126/sciadv.adv9451.
18. Simonson, T. S. (2015). Altitude adaptation: A glimpse through various lenses. High Altitude Medicine & Biology, 16(2), 125–137. https://doi.org/10.1089/ham.2015.0033 .
19. Majmundar, A. J., Wong, W. J., & Simon, M. C. (2010). Hypoxia-inducible factors and the response to hypoxic stress. Molecular Cell, 40(2), 294–309. https://doi.org/10.1016/j.molcel.2010.09.022 .
20. Schödel, J., & Ratcliffe, P. J. (2019). Mechanisms of hypoxia signalling: New implications for nephrology. Nature Reviews Nephrology, 15(10), 641–659. https://doi.org/10.1038/s41581-019-0182-z .