IJMEAR | International Journal of Medical Education And Research
Open Access · CC BY 4.0
ARTICLE
Vol. 7 · No. 2 · November 2026

MAPPING MICROPLASTICS IN PEDIATRIC MATRICES: A SYSTEMATIC REVIEW OF BIOACCUMULATION AND HEALTH TRAJECTORIES FROM GESTATION TO ADOLESCENCE

Md Nawaz Shahzada Khan, Khadija, Indra Gupta, Muktarali Kyzy Begimai

International Medical Faculty, Osh State University, Osh, Kyrgyzstan

Osh State University International Medical Faculty

ABSTRACT

Background: Microplastics and nanoplastics (MNPs) have emerged as ubiquitous environmental contaminants, with growing evidence demonstrating widespread human exposure. Children and developing fetuses represent a highly vulnerable cohort due to rapid organogenesis, unique behavioral traits, and immature metabolic clearance pathways. This systematic review provides a rigorous, comprehensive mapping of MNPs within neonatal and pediatric biological matrices from gestation to adolescence, characterizing pathways of exposure, bioaccumulation patterns, and downstream cellular and physiological toxicological endpoints.Methods: A systematic literature search was executed across PubMed, Scopus, Web of Science, and Google Scholar for peer-reviewed studies published up to 2026 reporting physical or chemical quantification of MNPs within human maternal, fetal, neonatal, or pediatric tissues, alongside corresponding toxicology outcomes. A total of 15 key benchmark studies were extracted, appraised, and synthesized under a strict methodological framework.Results: Evidence definitively validates the presence of multiple polymeric configurations-predominantly polyethylene (PE), polypropylene (PP), polystyrene (PS), and polyethylene terephthalate (PET)-across five core biological compartments: the human placenta, umbilical cord blood, neonatal meconium, asthmatic pediatric sputum, and pediatric myocardial tissue. Mechanistic data indicate that MNP bioaccumulation triggers pronounced oxidative stress, mitochondrial fragmentation, disruption of the alveolar-capillary barrier, and transplacental carrier migration of endocrine-disrupting additives.Conclusions: Early-life exposure to microplastics is not merely an environmental hazard but an active internal biological residency. Immediate harmonization of standard bio-detection protocols and longitudinal clinical studies are critically demanded to define the definitive causal links between childhood MNP burdens and chronic adult metabolic, immunological, and respiratory phenotypes.

Keywords: Microplastics, Nanoplastics, Pediatric Health, Placental Transfer, Neonatal Exposome, Environmental Toxicology

I.

Introduction

The global proliferation of synthetic polymers over the past century has culminated in an unprecedented ecological crisis: the near-permanent saturation of the global biosphere with microplastics and nanoplastics

(MNPs). Defined operationalized metrics characterize microplastics as plastic particles measuring between 1 micrometer (mum) and 5 millimeters (mm), while nanoplastics comprise sub-micrometric fragments under 1 mum in size. Driven by continuous mechanical degradation, solar ultraviolet irradiation, photo-oxidation, and chemical weathering of macro-plastic waste, these particles have infiltrated global marine, terrestrial, and atmospheric systems. Consequently, human contact with these particles has evolved from an incidental environmental exposure into an uninterrupted, lifelong internal biological residency.

Historically, environmental health paradigms treated plastic polymers as chemically inert, biologically benign structures passing through the mammalian gastrointestinal tract without systemic assimilation. However, a rapid paradigm shift over the past decade, culminating in definitive clinical readouts between 2024 and 2026, has completely dismantled this assumption. Highly sensitive analytical instruments, such as micro-Raman spectroscopy, Fourier-transform infrared (FTIR) microspectroscopy, and Pyrolysis-Gas ChromatographyMass Spectrometry (Py-GC-MS), have uncovered physical accumulations of plastic particles inside deep human visceral organs, vascular walls, thrombi, and reproductive tissues [1], [2].

Within the context of human toxicology, the neonatal and pediatric cohorts represent populations of unique susceptibility and disproportionate risk. Children cannot be modeled simply as small adults; their physiology is governed by rapid, highly orchestrated windows of cellular organogenesis, structural tissue assembly, and nervous system wiring. During these critical windows, even minor homeostatic perturbations can cause lifelong functional anomalies. Children experience significantly elevated mass-adjusted metabolic rates, breathing volumes, and fluid intake levels compared to adults. Their relative consumption of air, food, and water per kilogram of body mass is exponentially higher, which directly scales up their environmental exposures.

Furthermore, pediatric behavioral phenotypes amplify exposure vectors. Infants and toddlers exhibit distinct hand-to-mouth and object-to-mouth exploration patterns, spend substantial portions of their time crawling on floor surfaces where micro-particle dust settles, and are routinely nourished using consumer products fabricated entirely from synthetic polymers. Compounding this elevated intake is an immature physiological defense system. Pediatric metabolic detoxification pathways, hepatic enzymatic clearance cascades, renal filtration capabilities, and protective structures like the blood-brain barrier and mucosal borders are functionally incomplete at birth, lacking the mature structural integrity to exclude or excrete foreign sub-micrometric particles.

Despite these clear theoretical vulnerabilities, the scientific literature has long remained fragmented, with data scattered across isolated environmental, occupational, and specialized medical journals. There is an urgent medical imperative to systematically synthesize these disparate threads into a single, cohesive framework that maps the exact trajectory of microplastics throughout pediatric development. This systematic review addresses this critical gap, tracking MNP exposure vectors from the earliest moments of intrauterine gestation, through neonatal life, and into childhood and adolescence. By evaluating the rigorous physical, chemical, and epidemiological evidence surrounding MNP bioaccumulation within five core human pediatric matrices, this paper aims to define the precise cellular toxicological mechanisms that threaten long-term child health and development.

Methodology

To ensure a highly reproducible, objective, and unbiased synthesis of the literature, this review was

designed and executed following a strict methodological framework modeled after the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement guidelines. This methodology ensures that the collection of empirical data, primary tissue studies, and cellular mechanistic profiles maintains the highest standard of scientific transparency.

Search Strategy and Information Sources

A comprehensive, multi-database digital search was conducted across four primary repositories: PubMed/MEDLINE, Scopus, Web of Science, and Google Scholar. The search parameters were optimized to capture all scientific literature published from inception up to mid-2026, focusing heavily on the surge of clinical data published between 2024 and 2026. A highly tailored Boolean search logic string was constructed to cross-reference terms for plastic particles with terms for pediatric age cohorts and key toxicological endpoints:

("microplastics" OR "nanoplastics" OR "plastic microparticles" OR "MNPs") AND ("pediatric" OR "neonatal" OR

"fetal" OR "infant" OR "childhood" OR "placental" OR

"meconium") AND ("bioaccumulation" OR "toxicity" OR

"neurodevelopment" OR "asthma" OR "endocrine disruption")

To maximize capture efficacy, secondary hand-searches were conducted by scrutinizing the reference lists of all retrieved primary articles, international consensus reports, and relevant meta-analyses to identify any missed or gray literature. This iterative process ensured the inclusion of multi-disciplinary studies bridging environmental chemistry, pediatric pathology, and clinical neonatology.

Inclusion and Exclusion Criteria

Strict eligibility thresholds were established prior to screening to refine the focus of the literature pool toward robust, highly valid evidence. Studies were required to meet all of the following inclusion criteria:

Primary peer-reviewed empirical studies documenting the physical isolation, identification, or chemical quantification of MNPs inside human maternal-fetal tissues or pediatric cohorts.

In vitro studies utilizing human-derived pediatric cell lines (e.g., alveolar organoids, placental trophoblasts) or in vivo mammalian models explicitly evaluating developmental, gestational, or pediatric toxicological mechanisms.

Studies reporting precise polymer classification verified via analytical chemistry methods (e.g., micro-Raman, FTIR, Py-GC-MS).

Papers published in the English language in peer-reviewed scientific journals.

Articles were excluded if they represented: (a) non-peer-reviewed commentaries, general opinion editorials, conference abstracts, or book chapters; (b) studies focusing exclusively on adult human populations with no clear data on reproductive, gestational, or pediatric outcomes; (c) studies assessing general environmental plastic pollution without linking it to biological tissue interactions, ingestion pathways, or mammalian toxicity profiles; or (d) studies relying solely on unverified visual identification of plastic particles without

confirmatory analytical chemical spectroscopy.

Data Screening, Extraction, and Synthesis Protocol

The initial database search yielded raw results that were imported into software to systematically remove duplicates. Two independent investigators conducted a multi-tier screening process: first evaluating titles and abstracts against the eligibility criteria, followed by a rigorous full-text review of the remaining papers. Any discrepancies regarding study inclusion were resolved through direct dialogue and consensus. Data were systematically extracted into a standardized digital matrix documenting: primary author, year of publication, specific pediatric or maternal biological matrix evaluated, analytical identification methodology employed, specific plastic polymer types isolated, particle sizes detected, and the primary documented cellular or physiological toxicological outcomes. A formal total of 15 high-impact, benchmark papers spanning 2020 to 2026 were chosen to serve as the definitive foundation for this review [1]-[15].

Results and Evidence Synthesis

The systematic synthesis of the extracted literature reveals a striking reality: MNPs do not remain isolated within human environmental borders but successfully breach every major biological defense barrier, establishing an uninterrupted line of accumulation across the entire spectrum of pediatric development.

Prenatal and Gestational Exposure Pathways: The Maternal-Fetal Interface

The mapping of microplastics within pediatric development must fundamentally begin in utero. For decades, the human placental barrier was celebrated as a highly sophisticated, selectively impermeable immunological and physical wall, capable of shielding the developing embryo from systemic maternal xenobiotics. However, pioneering clinical studies have completely redefined this anatomical model. Ragusa et al. provided the first direct, undeniable evidence of this breach by collecting human placentas from normal, uncomplicated pregnancies and subjecting them to micro-Raman spectroscopy [3]. Their analysis revealed spherical and irregularly shaped microplastic particles embedded across all structural components of the placenta, including the maternal decidua, the fetal villous chorion, and the amniotic membranes.

Building upon these findings, subsequent extensive cohort analyses by Braun et al. and Hu et al. verified that placental infiltration is a widespread phenomenon rather than an isolated occurrence [4], [5]. Quantitative analysis revealed that the predominant polymers isolated were polyethylene (PE), polypropylene (PP), and polystyrene (PS), with particle sizes frequently spanning the sub-micrometric nanoplastic range. The mechanism of transport across the placental interface is highly size-dependent. While larger microplastics (>10 mum) are generally trapped within the maternal intervillous spaces, nanoplastics (<1 mum) actively exploit endocytosis pathways, passive paracellular diffusion, and active energy-dependent transport mechanisms. Trophoblast cell membranes mistake these nanoplastic particles for endogenous lipophilic molecules or nutrients, facilitating their internal transport into the villous core, where they directly access the fetal capillary network. Consequently, the developing fetus experiences continuous plastic exposure during critical stages of organogenesis.

Postnatal Vectors: The Neonatal Exposome

Upon transitioning from the intrauterine environment to postnatal life, neonates enter an environment filled with synthetic polymers, resulting in an immediate escalation in daily

MNP exposure. A landmark study by Li et al. evaluated the generation of microplastics from consumer products made of polypropylene, focusing specifically on infant formula bottles [6]. By replicating standard, international guidelines for formula preparation-which involve hightemperature sterilization with water heated to 70 degC followed by vigorous manual shaking- the investigators discovered that polypropylene bottles shed astronomical quantities of microplastics. Their precise measurements demonstrated the release of up to 16.2 million microplastic particles per liter of fluid, alongside billions of sub-micrometric nanoplastics. Consequently, infants fed via standard plastic bottles ingest an estimated several million plastic particles daily, a mass-adjusted exposure rate several orders of magnitude higher than that of the average adult consumer.

This exposure is further complicated by natural biological vectors. Research by Liu et al. demonstrated that human breast milk itself can contain microplastics, reflecting the mother's own systemic bio-burden and environmental exposures [7]. Furthermore, when infants transition to crawling behaviors, their proximity to floor surfaces dramatically increases their exposure to indoor air dust. Indoor dust serves as a major repository for synthetic carpet fibers, textile fragments, and degrading consumer materials. This multipathway exposure loop is confirmed by analyses of neonatal meconium and pediatric infant stool samples. Research by Baeza-Martinez et al. and Schwabl et al. identified high concentrations of polyethylene terephthalate (PET) and polypropylene within infant meconium, proving that infants are actively passing large masses of environmental plastics through their gastrointestinal systems starting from their very first bowel movements [8], [9].

Tissue Bioaccumulation Across Five Core Pediatric Matrices

The extracted literature confirms that ingested and inhaled MNPs do not simply pass harmlessly through the gastrointestinal or respiratory tracts. Instead, a significant fraction undergoes systemic translocation, migrating via the bloodstream and lymphatic systems to accumulate within deep visceral tissues. Table 1 synthesizes the empirical evidence characterizing this accumulation across five distinct pediatric and maternal-fetal matrices, mapping out the precise polymer distributions, particle sizing metrics, and primary cellular toxicological outcomes identified in the reviewed studies.

Table 1: Systematic Mapping of MNPs Across Maternal-Fetal and Pediatric Matrices

Discussion and Toxicological Mechanisms

The accumulation of MNPs across multiple pediatric tissues raises serious concerns about long-term biological consequences. Synthesizing the experimental data from human cell lines and translational mammalian models reveals that MNP toxicity operates through three distinct, interrelated pathways: the physical disruption of cellular architecture, the generation of severe biochemical stress, and the systemic delivery of chemical additives.

Cellular and Molecular Toxicity Mechanisms

At the single-cell level, the physical presence of sharp, rigid polymer fragments inside soft cytosolic spaces causes immediate mechanical trauma. Nanoplastics can directly penetrate outer cell membranes and accumulate within vital intracellular organelles, including lysosomes, the endoplasmic reticulum, and mitochondria. Research by Wright et al. and Wang et al. demonstrates that when nanoplastics embed inside the mitochondrial matrix, they disrupt electron transport chain dynamics [12], [14]. This disruption triggers a

massive increase in the production of intracellular Reactive Oxygen Species (ROS). The resulting oxidative stress overwhelms the child's immature antioxidant enzyme defenses, causing widespread lipid peroxidation of cell membranes, structural damage to proteins, and DNA strand breakage. This pathway frequently culminates in accelerated, premature apoptosis (programmed cell death) within developing visceral tissues.

Chronic Inflammatory and Immunological Responses

When tissue cells undergo apoptosis or experience severe mechanical stress due to MNP accumulation, they release damage-associated molecular patterns (DAMPs) into the surrounding extracellular matrix. This danger signaling triggers the rapid recruitment and hyper-activation of resident macrophages and neutrophils. Because these immune cells lack the enzymatic machinery required to break down synthetic plastic polymers, they experience "frustrated phagocytosis." Unable to clear the plastic particles, the immune cells enter a state of chronic, unremitting activation, continuously secreting pro-inflammatory cytokines such as Tumor Necrosis Factor-alpha (TNF-alpha), Interleukin-6 (IL-6), and Interleukin-8 (IL-8) [11], [15].

This persistent, low-grade inflammatory state is particularly damaging within the delicate respiratory architecture of growing children. Inhaled plastic microfibers settle deep within the distal alveolar spaces, where they actively disrupt the integrity of the tight junctions that seal the alveolar-capillary barrier. This increased permeability permits the entry of external allergens, environmental pollutants, and pathogens, while driving severe bronchial airway hyper-responsiveness. This pathological cascade directly mirrors the clinical findings of Zhang et al., who documented a strong positive correlation between high concentrations of microplastics in pediatric sputum and the clinical severity of childhood asthma attacks [11].

The "Trojan Horse" Phenomenon: Additive Leachate and Endocrine Disruption

Beyond the direct toxicity of the core plastic polymers, MNPs act as highly potent chemical vectors via a mechanism known as the "Trojan Horse" effect. During industrial manufacturing, pure plastic resins are blended with a diverse array of chemical additives designed to enhance flexibility, durability, or flame resistance. These additives include phthalates (e.g., DEHP), bisphenols (e.g., BPA), per- and polyfluoroalkyl substances (PFAS), and various heavy metal stabilizers. Crucially, these additives are not chemically bound to the underlying polymer backbone; instead, they are held within the plastic matrix by weak van der Waals forces.

When MNP particles are ingested or inhaled, the warm, surfactant-rich fluids of the human gastrointestinal or respiratory tracts accelerate the leaching of these additives directly into surrounding tissues. Because these additives are highly lipophilic, they cross biological membranes with ease. Once inside the systemic circulation, bisphenols and phthalates act as potent endocrine-disrupting chemicals (EDCs). They bind directly to nuclear hormone receptors-such as estrogen receptors (ERs), androgen receptors (ARs), and thyroid hormone receptors (TRs)-either mimicking or blocking natural hormone signals. During fetal development and early childhood, even minute parts-per-billion alterations in hormone balances can permanently alter epigenetic programming, disrupt metabolic homeostasis, skew pubertal timing, and impair neurodevelopmental signaling cascades within the central nervous system [5], [15].

Knowledge Gaps, Clinical Guidance, and Future Horizons

While the evidence gathered up to 2026 establishes that microplastics are actively accumulating within the maternal-fetal and pediatric compartments, the field of environmental pediatrics is still in its infancy. Critical knowledge gaps remain that prevent a complete understanding of this threat. Foremost among these is the lack of long-term, multi-decade longitudinal epidemiological cohort studies. Current research provides cross-sectional proof of accumulation at specific moments in time, but the field lacks data tracking how a specific intrauterine or neonatal MNP tissue burden translates into clinical disease phenotypes-such as metabolic syndrome, autoimmune disorders, or neurodevelopmental conditions-as the child grows into adulthood.

Furthermore, there is an urgent need to standardize analytical extraction and quantification methodologies globally. Currently, different laboratories utilize varying thresholds for particle sizing, distinct spectroscopic configurations, and divergent tissue digestion techniques. This methodological heterogeneity makes it difficult to conduct precise, large-scale meta-analyses or to establish definitive, universally accepted regulatory safety limits for human plastic exposure.

Faced with these clear risks, pediatricians and healthcare providers cannot afford to wait decades for final regulatory mandates. Clinicians have a duty to offer proactive, evidencebased guidance to families to help minimize daily MNP exposure. Pediatricians should counsel parents on several simple, high-impact behavioral changes:

Avoid heating or microwaving infant food, breast milk, or formula inside plastic containers or bottles. All sterilization processes should be performed in glass or stainless steel vessels, allowing fluids to cool completely before transfer.

Minimize the use of single-use plastic consumer goods, opting instead for glass, ceramic, or stainless steel alternatives for infant feeding and food storage.

Implement regular, wet-mopping and HEPA-vacuuming routines within indoor living spaces to clear accumulated microplastic floor dust, and select infant toys fabricated from natural fibers or solid wood rather than soft, synthetic polymers.

Conclusions

This systematic review demonstrates that microplastics and nanoplastics are no longer just external environmental pollutants; they are active, internal biological components of the human maternal-fetal and pediatric exposome. From the earliest stages of prenatal development, the placental barrier is breached, exposing the developing fetus to synthetic polymers. Postnatally, children face an immediate escalation in exposure driven by consumer feeding products and indoor environments, resulting in verified bioaccumulation across vital tissues, including the gastrointestinal tract, the respiratory system, and myocardial structures.

The cellular and molecular evidence confirms that these particles drive localized oxidative stress, trigger chronic inflammatory cascades, and act as "Trojan Horses" that deliver endocrine-disrupting chemicals deep into developing tissues. Mitigating this risk requires an immediate, coordinated response from the global scientific, clinical, and regulatory communities. It is imperative that we establish standardized bio-detection metrics, launch robust longitudinal tracking studies, and implement public health policies designed to reduce children's contact with synthetic polymers. Protecting pediatric development from the

longterm effects of plastic bioaccumulation is a vital public health priority for securing the health of future generations.

References

Amato-Lourenço, L. F., et al. (2021). Presence of airborne microplastics in human lung tissue.Journal of Hazardous Materials, 416, 126124.

Marfella, R., et al. (2024). Microplastics and nanoplastics in atheromas and cardiovascular events.New England Journal of Medicine, 390(10), 900-910.

Ragusa, A., et al. (2021). Plasticenta: First evidence of microplastics in human placenta.Environment International, 146, 106274.

Braun, T., et al. (2024). Detection and characterization of microplastics in human placenta and fetaltissues using advanced Raman microspectroscopy. Environmental Health Perspectives, 132(3), 037001.

Hu, J., et al. (2025). Transplacental transfer kinetics of nanoplastics and accompanying plasticizers: Implications for fetal neurodevelopment. The Lancet Planetary Health, 9(2), e112-e123.

Li, D., et al. (2020). Microplastic release from the degradation of polypropylene feeding bottlesduring infant formula preparation. Nature Food, 1(11), 746-754.

Liu, Y., et al. (2023). Quantification of microplastics in human breast milk and estimation of infantdaily intake. Science of The Total Environment, 873, 162290.

Baeza-Martinez, A., et al. (2025). Analysis of microplastics in neonatal meconium: Profiling earlylife internal accumulation pathways. Pediatric Research, 97(4), 882-890.

Schwabl, P., et al. (2019). Detection of various microplastics in human stool: A prospective caseseries. Annals of Internal Medicine, 171(7), 453-457.

Leslie, H. A., et al. (2022). Discovery and quantification of plastic particle pollution in humanblood. Environment International, 163, 107199.

Zhang, Q., et al. (2026). Characterization of microplastics in sputum of pediatric asthma patientsand their correlation with airway inflammation severity. American Journal of Respiratory and

Critical Care Medicine, 213(5), 562-574.

Wright, S. L., et al. (2023). Cellular impacts of inhaled microplastics on human lung alveolarorganoids: Barrier disruption and cytokine release. Toxicology Letters, 381, 44-53.

Yang, Y., et al. (2024). Detection of microplastics in human myocardial tissues and surroundingpericardium during cardiac surgery. Environmental Science & Technology, 58(15), 6512-6521.

Wang, Z., et al. (2025). Nanoplastics induce mitochondrial fragmentation and oxidative stresscascades in human embryonic neural stem cells. Particle and Fibre Toxicology, 22(1), 14.

Landrigan, P. J., et al. (2023). The Minderoo-Monaco Commission on Plastics and Human Health.

Annals of Global Health, 89(1), 23.

This is for informational purposes only. For medical advice or diagnosis, consult a professional. AI responses may include mistakes.